



Working together to improve health care quality, outcomes, affordability, and equity in Washington State

Pediatric Asthma

Draft 2022

Executive Summary

Asthma is the most common chronic disease among children,¹ with many potential risk factors including smoking, obesity, family history, and environmental triggers, or adverse childhood experiences.² From 2008 to 2013 the estimated annual economic cost of asthma was \$81.9 billion due to medical costs, loss of work and school days, and asthma-related mortality.³ In Washington state nearly 120,000 children have been diagnosed with asthma, accounting for between 8 to 11 percent of children in middle and high school.⁴ Additionally, the burden of asthma in the United States falls disproportionately on Black, Hispanic, and American Indian/Alaskan Native people.⁵ Asthma cannot be cured but can be effectively managed through strong treatment and mitigation plans.

The Bree Collaborative elected to develop recommendations for pediatric asthma in 2022 and convened a workgroup of subject matter experts from January to November 2022. Pediatric asthma control requires collaboration across numerous partners ranging from pediatricians to school nurses. In response, the Bree Collaborative's pediatric asthma workgroup includes providers, plans, public health agencies, patient advocates, and school representatives. This guideline is meant to address asthma triggers and control across numerous sites and ensure communication across sectors.

Recommendations focus on asthma control in the **clinical environment, home environment, school environment, and community environment**. Additionally, recommendations are directed toward:

- Health systems leadership
- Clinicians
- Home-based interventions
- Schools
- Public health agencies
- Those receiving care (patients/consumers)

We summarize potential avenues to improve alignment for pediatric asthma care through a review of published evidence, best practices from existing interventions, and subject matter expert collaboration across health, public, and social sectors.

We hope this guideline increases collaboration across programs where children with asthma receive care and improves effective asthma management for all children in Washington state.

Bree Collaborative Pediatric Asthma Guidelines

Updated: September 13, 2022

Bree Collaborative Background

Washington State House Bill 1311 established the Dr. Robert Bree Collaborative in 2011 “...to provide a mechanism through which public and private health care stakeholders can work together to improve quality, health outcomes, and cost effectiveness of care in Washington State.” The Bree Collaborative was named in memory of Dr. Robert Bree, a leader in the imaging field and a key member of previous health care quality improvement collaborative projects.

Members are appointed by the Washington State Governor and include public health care purchasers for Washington State, private health care purchasers (employers and union trusts), health plans, physicians and other health care providers, hospitals, and quality improvement organizations. The Bree Collaborative is charged with identifying health care services annually with substantial variation in practice patterns, high utilization trends in Washington State, or patient safety issues. For each health care service, the Bree Collaborative identifies and recommends best-practice, evidence-based approaches that build upon existing efforts and quality improvement activities to decrease variation. In the bill, the legislature does not authorize agreements among competing health care providers or health carriers as to the price or specific level of reimbursement for health care services. Furthermore, it is not the intent of the legislature to mandate payment or coverage decisions by private health care purchasers or carriers.

See **Appendix A** for a list of current Bree Collaborative members.

Recommendations are sent to the Washington State Health Care Authority for review and approval. The Health Care Authority (HCA) oversees Washington State’s largest health care purchasers, Medicaid, and the Public Employees Benefits Board Program, as well as other programs. The HCA uses the recommendations to guide state purchasing for these programs. The Bree Collaborative also strives to develop recommendations to improve patient health, health care service quality, and the affordability of health care for the private sector but does not have the authority to mandate implementation of recommendations.

For more information about the Bree Collaborative, please visit: www.breecollaborative.org.

Bree Collaborative members identified pediatric asthma as a priority improvement area and convened a workgroup to develop evidence-informed standards. The workgroup met from January to November 2022.

See **Appendix B** for the workgroup charter and a list of members.

Background

Asthma is a long-term disease that causes inflammation and swelling of the airways. Often, people with asthma are sensitive to “triggers” that may not bother other people, which can include dust mites, pollen, molds, cigarette smoke, or air pollution.⁶ About 1 in 12, or 6 million children in the US ages 0 – 17 have asthma,⁷ the highest asthma prevalence rate of any other age group in 2018.⁸ Additionally, it is estimated that 50% of children with asthma have uncontrolled asthma.⁹ In the United States, the burden of asthma falls disproportionately on Black, Hispanic, and American Indian and Alaskan native populations with disparities present in asthma prevalence, mortality, and health care utilization.¹⁰ The estimated annual economic burden of asthma in the United States is \$81.9 billion.¹¹

Washington state has been identified as having one of the highest asthma prevalence in the nation, with more than half a million adults and nearly 120,000 youth currently diagnosed with asthma.¹² Over one year in Washington state, 28% of youth with asthma lacked routine healthcare visits, and one in five youth with asthma visited an emergency room to manage asthma symptoms.¹³ Additionally, poor control of pediatric asthma can lead to chronic absenteeism and affect the academic success of children.¹⁴

While we don’t know exactly what causes asthma, we can prevent asthma attacks or at least make them less severe.¹⁵ Effectively managing pediatric asthma will require collaboration across multiple stakeholders who care for children, including pediatricians, school nurses, and public health agencies.

Given the multidisciplinary nature of pediatric asthma control, this guideline focuses on strategies to align efforts across key stakeholders. Recommendations are meant to supplement existing treatment guidelines from the [National Asthma Education and Prevention Program](#),¹⁶ best practices for community interventions from the [CDC’s Community Preventative Services Task Force](#),¹⁷ and expectations for pediatric asthma care in schools from [OSPI’s Asthma Management in Educational Settings](#) guide.¹⁸ Additionally, this guideline builds off previous work to mitigate the effects of climate on asthma from the [Asthma and Allergy Foundation of America](#)¹⁹ and offers strategies for funding pediatric asthma interventions drawing from previous research on funding mechanisms from the [Brookings Center for Health Policy](#).²⁰ The Bree workgroup’s focus areas are detailed in **Table 1**.

Table 1: Bree Collaborative Pediatric Asthma Focus Areas

Focus Areas	Goal(s)
Clinical Control	<ul style="list-style-type: none">• Appropriately establish asthma diagnosis and assess for severity and risk.• Develop an asthma management plan to effectively control asthma.• Implement appropriate asthma metrics to ensure quality care.
Home Environment	<ul style="list-style-type: none">• Define home-based interventions to address pediatric asthma.• Offer recommendations to operate and scale up home-based interventions.

Bree Collaborative Pediatric Asthma Guidelines

Updated: September 13, 2022

	<ul style="list-style-type: none">• Recommend potential sources of sustainable funding for home-based interventions.
School Environment	<ul style="list-style-type: none">• Appropriately manage pediatric asthma in the school setting.• Improve communication between school nurses, school-based health centers, and pediatricians/clinical providers.
Community Environment/Climate	<ul style="list-style-type: none">• Mitigate the effects of climate change, air pollution, and other environmental triggers on pediatric asthma.• Develop strategies to respond to environmental triggers
Funding	<ul style="list-style-type: none">• Consider alternative funding models for pediatric asthma that prioritize prevention and control.

Recommendations

Recommendations for Health Delivery Systems

Clinical Environment

- Ensure asthma management policies and procedures follow the most recent NHLB guidelines.
- Implement automated notification systems for asthma diagnosis.
 - a. Implement automated notification system when a patient is diagnosed with asthma when admitted to the emergency room, urgent care, or inpatient hospitalization.
- Develop metrics for asthma control quality.
 - a. Track and measure asthma prevalence and stratify by severity and control.
 - i. Track severity and control using ICD-10 codes.
 - b. Measure the Asthma Medication Ratio to track population health or alternatively measure the frequency of bronchodilator prescription refills.
 - c. Measure rates of annual flu shot administration among children with asthma.

Community Environment/Climate

- Review [Climate Action Handbook for Hospitals](#) and consider making policy changes.
- Aim for carbon neutrality by reducing carbon emissions through renewable energy, greener care delivery, and low- and zero-emission transportation.
- Plan for climate mitigation infrastructure including adequate air filtration. Monitor clinical air quality.
- Enforce smoke-free policies
- Discuss air pollution mitigation when educating patient and family members about asthma management.
- Use environmentally-friendly, EPA-approved disinfectants for cleaning

Recommendations for Clinicians

Clinical Environment

- Appropriately establish an asthma diagnosis according to NHLB Guidelines
 - Determine that signs and symptoms of recurrent reversible airway obstruction are present based on history and exam. Consider other causes of obstruction.
 - For children over 5 years old: In addition to patient history, use spirometry to determine lung function. Rule out other potential conditions that can cause asthma-like symptoms such as respiratory tract infections.
 - In absence of other diagnostic tests (spirometry), determine if symptoms respond to albuterol.
 - For children under 5 years old: rely on information from the patient and caregiver. Diagnostic tests are not accurate for children under 5.
 - If the asthma appears to be triggered by environmental allergies, recommend allergy skin testing or blood testing for allergies.

Bree Collaborative Pediatric Asthma Guidelines

Updated: September 13, 2022

- Consider referral to a specialist (pediatric pulmonologist, allergist, asthma specialist) if diagnosis or testing is unclear.
- Assess for asthma severity (impairment and risk) and control according to most recent NHLB guidelines.
 - Assess for severity at initial visit and reassess severity at least annually. Include assessment for comorbid conditions and/or environmental exposures.
 - Assess for control at all planned asthma visits, at least annually. Include assessment of proper medication administration technique, suitability of asthma action plan, patient adherence, and patient/caregiver concerns.
- Develop an asthma management plan that includes exposure mitigation, medication, and immunotherapy as needed.
 - Implement mitigation interventions for asthma triggers related to indoor allergens (e.g. dust mites), outdoor allergens (e.g. pollen), and/or irritants (e.g. smoke). Ensure access to smoking cessation interventions to reduce exposure to second-hand smoke.
 - Implement a medication plan for asthma management.
 - Initiate corticosteroid and rescue treatment as appropriate. Consider dry powder inhalers (DPIs) for older children and teens who can tolerate DPIs.
 - Follow the 2020 updates to NHLB guidelines for treatment using inhaled corticosteroids (ICS) and short-acting beta₂-agonists (SABA)
 - Refer to a specialist to provide immunotherapy if required.
 - For individuals ages 5 years and older with mild to moderate allergic asthma, recommend use of subcutaneous immunotherapy as an adjunct treatment to standard pharmacotherapy.
 - For individuals with persistent allergic asthma, recommend against the use of sublingual immunotherapy.
- Schedule planned preventative visits for asthma control at least annually.
 - Ensure the asthma management plan reduces impairment and risk. Implement the Asthma Control Test to determine the patient's level of asthma control at every visit.
 - Schedule additional planned preventative visits specific to asthma control as needed. Review and update the asthma control plan as needed. Normalize routine asthma control visits.
 - Routine asthma control visits should also include multifaceted approaches to patient education, control of environmental triggers, and/or comorbid conditions that affect asthma.
 - Telehealth visits may be appropriate for follow-up and asthma education.

Alignment and Communication

- Communicate asthma management plan with external partners and the broader care team including school nurses and community health workers.
- Establish clear lines of communication with the patient's school nurse. Ask the parent/caregivers to complete a release of information form to allow bi-directional communication about the asthma plan.
 - Send printed discharge instructions or other forms from the child's asthma control visits to the school nurse

Bree Collaborative Pediatric Asthma Guidelines

Updated: September 13, 2022

- Ask parents/caregivers to fill out release forms while at their asthma control visit
 - Work to standardize delivery system liability waivers to facilitate information sharing.
- Align educational efforts about inhaler use and asthma management with the school-based asthma care plan.
- Provide asthma patients with a prescription for an inhaler and a spacer, as well as a second inhaler and spacer for school use
 - Plans: cover a second inhaler and spacer for school use at the start of the school year

Recommendations for Home-Based Interventions and Community Health Workers

Home Environment

- Home-based multi-trigger, multicomponent interventions can reduce exposure to multiple indoor asthma triggers, including allergens and irritants. These interventions should include home visits by trained personnel to:
 - Assess the home environment
 - Encourage smoke-free environments
 - Change the indoor environment to reduce exposure to asthma triggers such as mold removal, pest management, allergen management, and air filters.
 - Provide trigger-abatement products such as bedding encasements, vacuums, or cleaning supplies.
 - Provide education about asthma triggers present in the home environment.
- Programs may also offer non-environmental activities:
 - Training, motivational interviewing, and goal setting to improve asthma self-management.
 - General asthma education on self-care including medication adherence, inhaler technique, symptom management, and trigger assessment/reduction
 - Social services and support, including coordinated care for the asthma client such as referrals to resources and legal/housing assistance.
 - Home-based interventions should consider implementing social need screening tools based on previous [Bree recommendations](#).
 - Combine asthma-related interventions with other health interventions, such as teaching lead-poisoning prevention and offering vaccinations.
- Home-based interventions should:
 - Address all components of the NHLBI asthma guidelines.
 - Consider behavioral and social drivers of asthma disparities.
 - Be tailored to diverse populations and programs, with expanded outreach for patients and families experiencing asthma disparities.
 - Include follow-up with multiple visits (at least 3) to build relationships with patients and caregivers. Interventions should include a face-to-face component for visits, whether in-person or through virtual means.
 - Hire and train community health workers to improve outreach.
 - Follow the [NCQA/Penn Medicine guidelines](#) for supporting community health workers.

Alignment and Communication:

Bree Collaborative Pediatric Asthma Guidelines

Updated: September 13, 2022

- Pediatric patients admitted to the hospital for asthma should be referred for a home-based assessment and intervention where available
- Consider implementing social needs screening tools in the clinical setting ([Bree recommendations](#))
- Link patient screening results to community resources – either with community resource lists, community information exchanges, MCOs, etc
- Form strong partnerships and communication policies between health and social service organizations.
- Invest in bidirectional care coordination solutions.
- Empower patients and family/caregivers to navigate communication with providers and plans.
- Open channels of communication between community health worker activities and primary care providers. Fill out a release of information as needed.
 - Send visit summaries about community health worker interventions to PCPs
 - Communicate patient needs to community health worker programs
- Connect CHWs with MCOs for case management, durable medical equipment, and medication changes.

Recommendations for Schools and School Nurses

School Environment

- Work to identify students with asthma
 - Monitor students for signs of asthma, including students with unusually high absence rates and multiple office visits for airway obstruction.
 - Solicit asthma plan forms from providers and include checkboxes for severity and control.
- Develop a care plan for all students with asthma and update the care plan at least annually.
 - Screen students for asthma control using the Asthma Control Test and refer to their primary care provider if asthma is not controlled.
 - Include education on proper inhaler technique and medication management as part of the care plan.
 - Refer to existing resources for managing asthma in the school environment, including the OSPI Asthma Management in education Settings (AMES) guideline, the AASA Addressing Childhood Asthma recommendations, and the CDC Strategies for Assessing Asthma in Schools.
- Communicate the student's care plan with parents/caregivers, the student's pediatrician or other clinical provider, and other school staff, including teachers.
 - Engage parents/caregivers about strategies for managing asthma. Ask parent/caregivers to complete a release of information form to allow bi-directional communication about the asthma plan.
 - Establish communication with the student's pediatrician or other clinical provider. Ask providers to send printed discharge instructions or other forms from the child's asthma control visits to the school nurse.
- Ensure healthy school environments for asthma management and control.

Bree Collaborative Pediatric Asthma Guidelines

Updated: September 13, 2022

- Follow RCW 28A.210.370 to ensure in-service training on asthma, adoption of policies for asthma rescue, and allow students to carry and self-administer prescribed asthma medication.
- Adopt strategies to improve indoor air quality to reduce asthma triggers, including proper classroom air filtration systems to address air pollution and reduce exposure to respiratory viruses.
 - Refer to [ASHRAE](#) for indoor air quality guidelines for schools
- Adopt strategies to address air pollution and outdoor smoke, including following the DOH's [Outdoor Air Quality and School Activities](#) guide, and ensuring a tobacco and vape free campus.
- Follow AAP guidelines on school nurse staffing – employ at least one school nurse per building. Involve school nurses in air quality control plans and health plans.
- If the school has a school-based health center, offer asthma management and education services to students enrolled with the school-based health center (SBHC).
 - SBHC providers refer to the recommendations for clinicians for additional guidance on diagnosing and managing asthma.
 - Offer to administer asthma control medication on-site.
 - Provide follow-up care to students following an urgent care visit related to asthma.
 - Communicate the student's asthma control plan with the school nurse and other provider as necessary. Coordinate logistics of SBHC with school nurses.
 - Work with school nurses to run risk screening and update asthma care plan.

Alignment and Communication:

- Establish clear lines of communication with the patient's pediatrician or regular healthcare provider. Ask the parent/caregivers to complete a release of information form to allow bi-directional communication about the asthma plan.
 - Request printed discharge instructions or other forms from the child's asthma control visits to the school nurse.
 - Ask parents/caregivers to fill out release forms before the beginning of the school year.
 - Work to standardize delivery system liability waivers to facilitate information sharing.

Recommendations for Public Health Agencies

School Environment:

- Consider expanding the school-based health center model to provide on-site medical services mindful of the school environment.
- Develop and update guidance for improving indoor (e.g. HVAC systems) and outdoor (e.g. source control) air quality in schools.
- Develop best practices for managing asthma at childcare centers.
- Advocate for legislation and funding to stock albuterol and spacers for school nurse offices, similar to current policy for epi pens.

Community Environment/Climate:

Bree Collaborative Pediatric Asthma Guidelines

Updated: September 13, 2022

- Provide education to the community about climate change, air pollution, and risks to health
- Develop preparedness and mitigation plans for extreme weather events – including wildfires, extreme heat/cold, and flooding.
- Offer programs to connect patients and caregivers to air filters and other climate mitigation products.
- Partner with weatherization programs from the [Department of Commerce](#) to provide products to improve air quality.
- Improve options for clean transportation including bikes, carpooling, zero-emission buses.
- Advocate for a climate lens in healthcare policy including funded programs and research to prepare for the health impacts of a changing climate.
- Remove requirement to use chlorine bleach as a disinfectant
- Enforce the [National Center for Health Housing's](#) recommendations for air quality codes for personal homes and facilities. Provide funding to encourage upgrades to meet air quality codes.

Recommendations for Health Plans

- Cover routine asthma control visits for patients with asthma.
- Provide coverage for corticosteroids and rescue treatment for asthma patients, including coverage dry powder inhalers.
- Provide coverage for annual flu shots for children with asthma.
- Measure the Asthma Medication Ratio or the frequency of bronchodilator prescription refills to track population health.
- Connect home insured patients with asthma to climate abatement products, including HEPA filters.
- Consider payment models other than fee-for-service, including alternative payment models such as bundled payments that will help offset costs for care coordination.

Recommendations for Patients and Community Members

- Manage exposure to air pollution, especially during times of high wildfire smoke.
- Consider purchasing an air filter or working with a community organization to receive a discounted air filter.
- Maintain a smoke-free environment in the home and in vehicles, including tobacco, marijuana, vaping. Minimize wood stove and fireplace use. Turn on fan whenever using a gas stove.

Review of Evidence

Clinical Environment

Clinical guidelines for asthma are well-established, including recommendations from the [Global Initiative for Asthma \(GINA\)](#) and the National Heart, Lung, and Blood Institute (NHLBI) out of the National Institutes of Health.^{21,22} The NHLBI's National Asthma Education and Prevention Program Coordinating Committee convened their fourth expert panel working group to provide focused updates to their asthma guidelines in 2020. The updated guidelines focus on six priority topics including the use of fractional exhaled nitric oxide diagnostic tests, remediation of indoor allergens, adjustable medication dosing, long acting antimuscarinic agents, immunotherapy, and bronchial thermoplasty.²³

Overall, the NHLBI guidelines offer steps to diagnose new asthma cases, assess asthma severity based on risk and impairment, develop asthma control plans including medication therapy, and planning for routine asthma control visits.²⁴ The NHLBI guidelines for asthma management are mostly the same for adults and pediatric populations, except for diagnosis. As children younger than 6 years cannot take a spirometry diagnostic test, the NHLBI recommends using a medical history and symptoms to confirm asthma diagnosis for this age group.²⁵

The main asthma quality metric required by the Healthcare Effectiveness Data and Information Set (HEDIS) is the asthma medication ratio (AMR), which assesses adults and children 5–64 years of age who were identified as having persistent asthma and had a ratio of controller medications to total asthma medications of 0.50 or greater during the measurement year.²⁶ One study of children on Medicaid found that the AMR reliably predicted asthma exacerbations.²⁷ Alternatively, this workgroup suggests measuring the frequency of bronchodilator prescription refills to track population health. Bronchodilators serve as rescue inhalers for treating sudden asthma symptoms. Measuring the frequency of bronchodilator prescription refills can help track patients who are using their rescue inhaler more than recommended.

This workgroup offers a few clinical recommendations that extend beyond the NHLBI. Specifically, this workgroup recommends appropriate telehealth visits for routine asthma control visits and asthma education, depending on the patient's level of asthma control and in line with the [Bree Collaborative's 2021 Telehealth Guidelines](#).²⁸ Additionally, this workgroup recommends making dry powder inhalers available for children and youth who have the lung capacity to tolerate them. Several studies have demonstrated that dry powder inhalers are as effective as pressurized meter dose inhalers in controlling asthma, but can reduce the carbon footprint by half.^{29,30}

Home Environment

Home-based multi-trigger, multicomponent interventions can reduce exposure to multiple indoor asthma triggers (allergens and irritants).³¹ These interventions involve home visits by trained personnel to:

- Assess the home environment and change the indoor environment to reduce exposure to asthma triggers through processes including mold removal, pest management, allergen management, encouraging smoke-free behaviors, and connecting families to air filters.
- Provide education about the home environment

Bree Collaborative Pediatric Asthma Guidelines

Updated: September 13, 2022

Home-based asthma interventions may also include other activities to address patient and caregivers' barriers to receiving asthma care. Additional activities may include:

- Training, motivational interviewing, and goal setting to improve asthma self-management
- General asthma education on self-care including medication adherence, inhaler technique, symptom management, and trigger assessment/reduction
- Social services and support, including coordinated care for the asthma client such as referrals to resources and legal/housing assistance
- Combine asthma-related interventions with other health interventions, such as teaching lead-poisoning prevention and offering vaccinations.

The Community Preventative Services Task Force (CPSTF), a program of the US Department of Health and Human Services strongly recommends home-based multi-trigger, multicomponent interventions as an evidence-based intervention to address asthma. More than 22 studies have demonstrated that home-based interventions reduce asthma symptom days, days of school missed, acute healthcare visits, and increase pulmonary function.³² Additionally, the Asthma and Allergy Foundation of America have identified home-based interventions as a key strategy for reducing asthma disparities, especially for low-income and BIPOC communities.³³ In Washington state, the Healthy Homes program out of Public Health of Seattle and King County has demonstrated that a home-based program supported by community health workers can reduce asthma morbidity.^{34,35}

Successful home-based interventions share several key components including face-to-face components (virtual or in-person), tailoring the program diverse populations, and addressing behavioral and social drivers of asthma disparities. NCOA/Penn Medicine offer additional guidelines for supporting successful community health worker interventions.³⁶ Home-based community health worker programs may also present the opportunity to address patient and caregiver's social needs. See the Bree Collaborative endorsed recommendations on addressing social needs and health equity for more information.³⁷

School Environment

Schools are an essential partner to help address pediatric asthma, as students are expected to enroll in school from preK to 12th grade and school nurses are expected to develop individualized health plans (IHPs) for students with asthma. Additionally, RCW 28A.210.370 outlines requirements for school districts to develop policies to train staff to manage asthma and develop plans to provide asthma rescue treatment for students as needed.

Beyond the requirements, several organizations have developed guidelines for managing asthma in the school setting. The Asthma Management in the Schools Task Force, convened by the American Lung Association, the Office of the Superintendent of Public Instruction, the Department of Health, and School Nurses in Washington state developed a resource guide for asthma management in educational settings.³⁸ Additionally, the CDC have developed strategies for addressing asthma in schools and the National Schools Board Association have developed steps to address childhood asthma in the community.^{39,40}

Full time school nurses are effective in improving student asthma management and reducing absenteeism.⁴¹ Standard operating procedure for school nurses managing asthma includes providing

Bree Collaborative Pediatric Asthma Guidelines

Updated: September 13, 2022

asthma education, developing asthma care plans, and addressing potential asthma triggers to the best of their ability. The Asthma and Allergy Foundation of America provides templates for asthma action plans that can be communicated between clinics and schools.⁴² One of the largest barriers to school-based asthma management is infrequent communication between clinics and schools.⁴³ This workgroup recommends increased communication and alignment across clinics and schools as a key component to improve holistic asthma management.

Beyond school nurses, school administration plays a role in improving asthma management in schools. School administration must provide training on asthma, adopt policies for asthma rescue care, and allow students to carry and self-administer prescribed asthma medication. In addition to what is required by the RCW, school administration can adopt strategies to reduce asthma triggers, including ensuring a tobacco and vape-free campus.⁴⁴ Additionally, school administration should refer to the Washington State Department of Health's (DOH) resources to address air pollution and outdoor smoke.⁴⁵ The DOH provides resources to guide appropriate activities during periods of poor outdoor air quality and smoke.⁴⁶ Finally, school administration can ensure each student has access to a school nurse by following AAP school nursing staffing guides and ensure each school site employs at least one full-time school nurse.⁴⁷

School-based health centers are another option for providing asthma management and education services to children in the school environment. School-based health centers are effective in reducing the risk of hospitalization and ED visits for children with asthma.⁴⁸ Washington state is home to more than 65 school based health centers in over 25 school districts.⁴⁹ School based health centers can help administer asthma control medication, provide follow-up care to students following an urgent care visit, and provide additional asthma management education as needed.

Community Environment and Climate

The community environment can exacerbate asthma symptoms depending on the potential triggers present in the area. The CDC, the AAFA, and the APHA warn that climate change in particular will increase the burden of air pollution, allergens, and wildfires, all of which can trigger asthma symptoms.^{50,51,52} Strong prevention and mitigation policies are needed to address the risk of climate change and the environment on asthma exacerbations.

The healthcare system in the US is responsible for 8.5% of total greenhouse gas emissions in the nation.⁵³ Many believe that it is important for healthcare to lead efforts to reduce emissions, partially because of the large share of emissions⁵⁴ and to improve healthcare resilience and equity.⁵⁵ To help healthcare organizations address climate change, the Health Care Climate Council offers a Climate Action Playbook for clinics and hospitals.⁵⁶

Programs that address air pollution, especially from wildfire smoke, will help improve asthma management. The Washington Department of Commerce offers a weatherization program for low-income homes that can help offset the cost of air filters to help manage wildfire smoke.⁵⁷ Policies that improve or regulate indoor air quality for schools, homes, or hospitals can also help manage asthma triggers.^{58,59} Prompt action on climate mitigation and adaption will be essential to manage asthma with changing environments.

Funding

Bree Collaborative Pediatric Asthma Guidelines

Updated: September 13, 2022


Effectively managing asthma requires a multisector that expands beyond the clinical walls. Funding mechanisms that address home-based interventions and encourage collaboration across healthcare, schools, and community sites are needed to further pediatric asthma control. The Brookings Institute highlighted several opportunities for innovative funding models in 2015 that include potential bundled payments or value based payments.^{60,61}

Draft

Next Steps and Alignment

Washington state has been identified as having one of the highest asthma prevalence in the nation, with more than half a million adults and nearly 120,000 youth currently diagnosed with asthma.⁶² While we don't know exactly what causes asthma, we can prevent asthma attacks or at least make them less severe.⁶³ In Washington, the largest gaps in effectively managing pediatric asthma include appropriate diagnosis of new cases and coordinating asthma care across multiple environments, including the clinical, home, and school environment. A model for communication across health care stakeholders is listed in **Figure 1**.

Figure 1: Healthcare Stakeholder Alignment on Pediatric Asthma



We hope this report supplements existing guidelines for pediatric asthma and outlines steps for future alignment across stakeholders. Each organization has a unique role to play to achieve control of pediatric asthma in Washington state. By working together to improve alignment across sectors we can reduce the burden of asthma in our state.

Bree Collaborative Pediatric Asthma Guidelines

Updated: September 13, 2022

Appendix A: Bree Collaborative Members

Member	Title	Organization
Susie Dade, MS	Patient Advocate	
David Dugdale, MD, MS	Medical Director, Value Based Care	University of Washington Medicine
Gary Franklin, MD, MPH	Medical Director	Washington State Department of Labor and Industries
Stuart Freed, MD	Chief Medical Officer	Confluence Health
Mark Haugen, MD	Family Medicine	Walla Walla Clinic
Darcy Jaffe, MN, ARNP, NE-BC, FACHE	Senior Vice President, Safety & Quality	Washington State Hospital Association
Sharon Eloranta	Medical Director, Performance Measurement and Care Transformation	Washington Health Alliance
Norifumi Kamo, MD, MPP	Internal Medicine	Virginia Mason Franciscan Health
Angie Sparks, MD	Chief Medical Officer, Community Plan	UnitedHealthcare
Wm. Richard Ludwig, MD	Chief Medical Officer, Accountable Care Organization	Providence Health and Services
Greg Marchand	Director, Benefits & Policy and Strategy	The Boeing Company
Kimberly Moore, MD	Associate Chief Medical Officer	Franciscan Health System
Carl Olden, MD	Family Physician	Pacific Crest Family Medicine, Yakima
Drew Oliveira, MD	Executive Medical Director	Regence BlueShield
Mary Kay O'Neill, MD, MBA	Partner	Mercer
Kevin Pieper, MD	Chief Medical Officer	Kadlac Medical Center
Susanne Quistgaard, MD	Medical Director, Provider Strategies	Premiera Blue Cross
John Robinson, MD, SM	Chief Medical Officer	First Choice Health
Jeanne Rupert, DO, PhD	Provider	The Everett Clinic
Hugh Straley, MD (Chair)	Retired	Medical Director, Group Health Cooperative; President, Group Health Physicians
Shawn West, MD	Medical Director	Embright, LLC
Judy Zerzan, MD, MPH	Chief Medical Officer	Washington State Health Care Authority

Appendix B: Pediatric Asthma Charter and Roster

The Bree Collaborative Pediatric Asthma Charter and Roster

Problem Statement

Asthma is the most common chronic disease among children, with many potential risk factors including smoking, obesity, family history, and environmental triggers, or adverse childhood experiences.⁶⁴ In Washington state nearly 120,000 children have been diagnosed with asthma, accounting for between 8 to 11 percent of children in middle and high school.⁶⁵ As asthma cannot be cured but can be effectively managed, there is a need for strong treatment and mitigation plans to prevent burden of disease among youth in Washington.

Aim

To increase evidence-informed screening, diagnosis, monitoring, and treatment for pediatric asthma to reduce the burden of disease in Washington state.

Purpose

To propose practical and evidence-informed recommendations to the full Bree Collaborative on reducing the burden of pediatric asthma including:

- Helping patients and families achieve asthma control.
- Appropriately diagnosing asthma cases and defining medically high-risk asthma.
- Improving access to and implementation of treatment protocols.
- Understanding and adapting treatment plans based on social determinants of health.
- Developing and adopting metrics for medication compliance and treatment process.
- Partnering with patients to ensure medication compliance and educate families.
- Engaging community support and broader public health programs.
- Addressing home, school, and community environments.
- Developing sustainable funding for pediatric asthma interventions.
- Improving care coordination between providers, plans, and community health supports.
- Disseminating and informing clinicians of ongoing evidence-based guidelines.

Duties & Functions

The workgroup will:

- Research evidence-informed and expert-opinion informed guidelines and best practices (emerging and established).
- Identify current barriers and future opportunities for implementing interventions.
- Consult relevant professional associations and other stakeholder organizations and subject matter experts for feedback, as appropriate.
- Maintain an equity lens while developing recommendations.
- Meet for approximately nine months, as needed.
- Provide updates at Bree Collaborative meetings.

Bree Collaborative Pediatric Asthma Guidelines

Updated: September 13, 2022

- Post draft report(s) on the Bree Collaborative website for public comment prior to sending report to the Bree Collaborative for approval and adoption.
- Present findings and recommendations in a report.
- Recommend data-driven and practical implementation strategies including metrics or a process for measurement.
- Create and oversee subsequent subgroups to help carry out the work, as needed.
- Revise this charter as necessary based on scope of work.

Structure

The workgroup will consist of individuals confirmed by Bree Collaborative members or appointed by the chair of the Bree Collaborative. The Bree Collaborative director and program coordinator will staff and provide management and support services for the workgroup.

Less than the full workgroup may convene to: gather and discuss information; conduct research; analyze relevant issues and facts; or draft recommendations for the deliberation of the full workgroup. A quorum shall be a simple majority and shall be required to accept and approve recommendations to send to the Bree Collaborative.

Meetings

The workgroup will hold meetings as necessary. Bree Collaborative staff will conduct meetings, arrange for the recording of each meeting, and distribute meeting agendas and other materials prior to each meeting. Additional workgroup members may be added at the discretion of the Bree Collaborative director.

Name	Title	Organization
Annie Hetzel, MSN, RN	School Health Services Consultant	Office of the Superintendent of Public Instruction
Brad Kramer, MPA	Asthma and Community Health Worker Program Manager	Public Health, Seattle & King County
Christopher Chen, MD	Associate Medical Director	WA Health Care Authority
David Ricker, MD	Pediatric Pulmonologist	Mary Bridge Children's
Doreen Kiss, MD	Pediatrician/Peds Population Health	University of Washington
Edith Shreckengast, MS	Clinical Health Coach – Registered Dietitian	Community Health Plan of Washington
Julee Christianson	Health Schools Washington Director	Office of the Superintendent of Public Instruction
Kate Hastings	Senior Policy Expert	Scientific Consulting Group
Kate Guzowski		Community Health of Central Washington
Katie Paul, MD, MPH	Physician	Kaiser Permanente
LuAnn Chen, MD, MHA, FAAFP	Senior Medical Director	Community Health Plan of Washington
Mark LaShell, MD	Allergist-Immunologist	Kaiser Permanente
Michael Dudas, MD, FAAP	Pediatrician	Virginia Mason Medical Center

Bree Collaborative Pediatric Asthma Guidelines

Updated: September 13, 2022

Vicki Kolios, MSHSA, CPHQ	Senior Program Director, Surgical COAP & Spine COAP	Foundation for Health Care Quality
---------------------------	--	---------------------------------------

Draft

Bree Collaborative Pediatric Asthma Guidelines

Updated: September 13, 2022

References

- ¹ Ferrante, G., & Grutta, S. 2018. The Burden of Pediatric Asthma. *Frontiers of Pediatrics*.
<https://doi.org/10.3389/fped.2018.00186>
- ² WA Department of Health. (2010). Children and Youth with Asthma: Washington State 2009-2010.
<https://www.doh.wa.gov/Portals/1/Documents/Pubs/345-305-ChildAsthmaFactSheet.pdf>
- ³ Nurmagambetov, T., Kuwahara, R., & Garbe, P. 2018. The Economic Burden of Asthma in the United States, 2008–2013. *Annals of the American Thoracic Society*, 15(3), 348–356. <https://doi.org/10.1513/annalsats.201703-259oc>
- ⁴ WA Department of Health. (2022). Asthma Data.
<https://www.doh.wa.gov/dataandstatisticalreports/diseasesandchronicconditions/asthmaata>
- ⁵ Asthma and Allergy Foundation of America. 2020. Asthma Disparities in America: A Roadmap to Reducing Burden on Racial and Ethnic Minorities. Aafa.org. Accessed June 2022. Available:
<https://www.aafa.org/media/2743/asthma-disparities-in-america-burden-on-racial-ethnic-minorities.pdf>
- ⁶ Asthma and Allergy Foundation of America. 2021. What Causes or Triggers Asthma? Accessed August 2022. Available: <https://www.aafa.org/asthma-triggers-causes/>
- ⁷ Centers for Disease Control and Prevention. 2018. Vital Signs: Asthma in Children. Accessed August 2022. Available: <https://www.cdc.gov/vitalsigns/childhood-asthma/index.html>
- ⁸ American Lung Association. 2022. Asthma Trends and Burden. Accessed August 2022. Available:
<https://www.lung.org/research/trends-in-lung-disease/asthma-trends-brief/trends-and-burden>
- ⁹ Centers for Disease Control and Prevention. 2019. Uncontrolled Asthma Among Children, 2012-2014. Accessed August 2022. Available: https://www.cdc.gov/asthma/asthma_stats/uncontrolled-asthma-children.htm
- ¹⁰ Asthma and Allergy Foundation of America. 2020. Asthma Disparities in America: A Roadmap to Reducing Burden on Racial and Ethnic Minorities. Accessed March 2022. Available:
<https://www.aafa.org/media/2743/asthma-disparities-in-america-burden-on-racial-ethnic-minorities.pdf>
- ¹¹ Nurmagambetov, T., Kuwahara R., & Garbe P. 2017. The cost of Asthma in the United States. *Annals of the American Thoracic Society*. Accessed August 2022. Available: <https://www.thoracic.org/about/newsroom/press-releases/resources/asthma-costs-in-us.pdf>
- ¹² Washington State Department of Health. 2020. Asthma Data. Accessed August 2022. Available:
<https://doh.wa.gov/data-statistical-reports/diseases-and-chronic-conditions/asthma-data>
- ¹³ Washington State Department of Health. 2013. The Burden of Asthma in Washington State. Accessed August 2022. Available: <https://doh.wa.gov/sites/default/files/legacy/Documents/Pubs//345-240-AsthmaBurdenRept13.pdf>
- ¹⁴ Washington State Office of the Superintendent of Public Instruction. 2013. Asthma Management in Educational Settings. Accessed June 2022. Available:
<https://www.k12.wa.us/sites/default/files/public/healthservices/pubdocs/asthmamannual.pdf>
- ¹⁵ Centers for Disease Control and Prevention. 2018. Vital Signs: Asthma in Children. Accessed August 2022. Available: <https://www.cdc.gov/vitalsigns/childhood-asthma/index.html>
- ¹⁶ National Health Lung and Blood Institute. 2022. National Asthma Education and Prevention Program. Accessed February 2022. Available: <https://www.nhlbi.nih.gov/science/national-asthma-education-and-prevention-program-naepp>
- ¹⁷ Centers for Disease Control and Prevention Community Preventative Services Taskforce. 2013. Asthma Control: Home-Based Multi-Trigger, Multicomponent Environmental Interventions for Children and Adolescents with Asthma. Accessed March 2022. Available: <https://www.thecommunityguide.org/sites/default/files/assets/Asthma-Home-Based-Children.pdf>
- ¹⁸ Washington State Office of the Superintendent of Public Instruction. 2013. Asthma Management in Educational Settings. Accessed June 2022. Available:
<https://www.k12.wa.us/sites/default/files/public/healthservices/pubdocs/asthmamannual.pdf>
- ¹⁹ Asthma and Allergy Foundation of America. 2021. Key Issues: Climate and Health. Accessed July 2022. Available:
<https://www.aafa.org/climate-and-health/>

Bree Collaborative Pediatric Asthma Guidelines

Updated: September 13, 2022

-
- ²⁰ Farmer S., Patel K., George M., McStay F., Hart R., McClellan M. 2015. A Case Study in Payment Reform to Support Optimal Pediatric Asthma Care. Center for Health Policy at Brookings. Accessed July 2022. Available: <https://www.brookings.edu/wp-content/uploads/2016/06/asthma-case-study.pdf>
- ²¹ Global Initiative for Asthma. 2022. 2022 GINA Main Report. Accessed August 2022. Available: <https://ginasthma.org/gina-reports/>
- ²² National Heart, Lung, and Blood Institute. 2021. Asthma Management Guidelines: Focused Updates 2020. Accessed March 2022. Available: <https://www.nhlbi.nih.gov/health-topics/asthma-management-guidelines-2020-updates>
- ²³ National Heart, Lung, and Blood Institute. 2021. Asthma Management Guidelines: Focused Updates 2020. Accessed March 2022. Available: <https://www.nhlbi.nih.gov/health-topics/asthma-management-guidelines-2020-updates>
- ²⁴ National Heart, Lung, and Blood Institute. 2022. Asthma for Health Professionals: Clinical Practice Guidelines. Accessed March 2022. Available: <https://www.nhlbi.nih.gov/health-topics/asthma-for-health-professionals>
- ²⁵ National Heart, Lung, and Blood Institute. 2022. Asthma in Children. Accessed September 2022. Available: <https://www.nhlbi.nih.gov/health/asthma/children>
- ²⁶ NCQA. 2022. Asthma Medication Ratio (AMR). Accessed August 2022. Available: <https://www.ncqa.org/hedis/measures/medication-management-for-people-with-asthma-and-asthma-medication-ratio/>
- ²⁷ Andrews L., Brinton D., Simpson K., & Simpson A. 2018. A Longitudinal Examination of the Asthma Medication Ratio in Children with Medicaid. Journal of Asthma. 57(10). <https://doi.org/10.1080/02770903.2019.1640727>
- ²⁸ Bree Collaborative. 2021. Telehealth Guidelines. Accessed September 2022. Available: <https://www.qualityhealth.org/bree/wp-content/uploads/sites/8/2021/09/Telehealth-Guidelines-Final.pdf>
- ²⁹ Woodcock A., Janson C., Rees J., Frith L., Lofdahl M., Moore A., Hedberg M., & Leather D. 2022. Effects of switching from a metered dose inhaler to a dry powder inhaler on climate emissions and asthma control: post-hoc analysis. Thorax. doi: 10.1136/thoraxjnl-2021-218088
- ³⁰ Steuer G., Prais D., Mussaffi H., Mei-Zahav M., Bar-On O., Levine H., Gendler Y., Blau H., & Stafler P. 2018. Inspiromatic-safety and efficacy study of a new generation dry powder inhaler in asthmatic children. Pediatric Pulmonology. <https://doi.org/10.1002/ppul.24077>
- ³¹ The Community Guide. 2008. Asthma: Home-Based Multi-Trigger, Multicomponent Environmental Interventions – Children and Adolescents with Asthma. Accessed June 2022. Available: <https://www.thecommunityguide.org/findings/asthma-home-based-multi-trigger-multicomponent-environmental-interventions-children-adolescents-asthma>
- ³² The Community Guide. 2008. Asthma: Home-Based Multi-Trigger, Multicomponent Environmental Interventions – Children and Adolescents with Asthma. Accessed June 2022. Available: <https://www.thecommunityguide.org/findings/asthma-home-based-multi-trigger-multicomponent-environmental-interventions-children-adolescents-asthma>
- ³³ Asthma and Allergy Foundation of America. 2022. Asthma Disparities in America. Accessed March 2022. Available: <https://www.aafa.org/asthma-disparities-burden-on-minorities.aspx>
- ³⁴ Krieger J., Takaro T., Song L., & Weaver, M. 2005. The Seattle-King County Healthy Homes Project: A Randomized, Controlled Trial of a Community Health Worker Intervention to Decrease Exposure to Indoor Asthma Triggers. American Journal of Public Health. 95(4): 652-659. doi: 10.2105/AJPH.2004.042994
- ³⁵ Elkuglia N., Crocker M., Stout J., Bolt K., Weiner B., & Kramer B. 2021. Development of an Asthma Home-Visit Training Program for Community Health Workers and Their Supervisors in Washington State. Frontiers of Public Health. <https://doi.org/10.3389/fpubh.2021.674843>
- ³⁶ NCQA and Penn Medicine. 2021. Critical Inputs for Successful Community Health Worker Programs. Accessed March 2022. Available: <https://www.ncqa.org/wp-content/uploads/2021/11/Critical-Inputs-for-Successful-CHW-Programs-White-Paper-November2021.pdf>
- ³⁷ Foundation for Health Care Quality. 2021. Social Determinants of Health. Accessed September 2022. Available: <https://www.qualityhealth.org/bree/wp-content/uploads/sites/8/2021/05/FHCQ-Bree-SDoH-Equity-Final.pdf>

Bree Collaborative Pediatric Asthma Guidelines

Updated: September 13, 2022

-
- ³⁸ Washington Asthma Initiative. 2013. Taking Asthma Care to Schools: Asthma Management in Educational Settings. Accessed April 2022. Available: http://www.waionline.org/AMES_1-18-2014_final2.pdf
- ³⁹ American Association of School Administrators and the National School Boards Association. In the Schoolyard and Beyond: Addressing Childhood Asthma in Your Community. Accessed April 2022. Available: [https://www.aasa.org/uploadedFiles/Childrens_Programs/Asthma/In-the-Schoolyard%20\(English\).pdf](https://www.aasa.org/uploadedFiles/Childrens_Programs/Asthma/In-the-Schoolyard%20(English).pdf)
- ⁴⁰ Centers for Disease Control and Prevention. 2017. Strategies for Addressing Asthma in Schools. Accessed April 2022. Available: https://www.cdc.gov/asthma/pdfs/strategies_for_addressing_asthma_in_schools_508.pdf
- ⁴¹ Rodriguez E., Rivera D., Perlroth D., Becker E., Wang N., & Landau M. 2013. School nurses' role in asthma management, school absenteeism, and cost savings: a demonstration project. J Sch Health. 83(12):842-50. doi: 10.1111/josh.12102.
- ⁴² Asthma and Allergy Foundation of America. 2021. Asthma Action Plan Templates. Accessed April 2022. Available: <https://www.aafa.org/asthma-treatment-action-plan/>
- ⁴³ Louisias M., Goldmann D., Phipatanakul W. 2016. School Nurses' Perspectives on Barriers to Implementing School-Based Asthma Management Plans. 137(2). <https://doi.org/10.1016/j.jaci.2015.12.453>
- ⁴⁴ American Association of School Administrators and the National School Boards Association. In the Schoolyard and Beyond: Addressing Childhood Asthma in Your Community. Accessed April 2022. Available: [https://www.aasa.org/uploadedFiles/Childrens_Programs/Asthma/In-the-Schoolyard%20\(English\).pdf](https://www.aasa.org/uploadedFiles/Childrens_Programs/Asthma/In-the-Schoolyard%20(English).pdf)
- ⁴⁵ Washington State Department of Health. 2022. Asthma and Schools. Accessed April 2022. Available: <https://doh.wa.gov/you-and-your-family/illness-and-disease-z/asthma/asthma-and-schools>
- ⁴⁶ Washington State Department of Health. 2022. Washington Air Quality Guide for School & Child Care Activities. Accessed April 2022. Available: <https://doh.wa.gov/sites/default/files/legacy/Documents/Pubs//334-332.pdf>
- ⁴⁷ American Academy of Pediatrics. 2016. Role of the School Nurse in Providing School Health Services. Pediatrics. 137(6). <https://doi.org/10.1542/peds.2016-0852>
- ⁴⁸ Guo J., Kang R., Keller K., McCracken A., Pan W., Cluxton R. 2005. Impact of school-based health centers on children with asthma. J Adolesc Health. 37(4): 266-74. doi: 10.1016/j.jadohealth.2004.09.006.
- ⁴⁹ Washington School-Based Health Alliance. 2022. SBHCs in Washington. Accessed September 2022. Available: <https://wasbha.org/sbhcs-in-washington-2/>
- ⁵⁰ Centers for Disease Control and Prevention. 2022. Climate Effects on Health. Accessed August 2022. Available: <https://www.cdc.gov/climateandhealth/effects/default.htm>
- ⁵¹ Asthma and Allergy Foundation of America. 2021. Climate and Health. Accessed August 2022. Available: <https://www.aafa.org/climate-and-health/>
- ⁵² American Public Health Association. 2021. Climate Change. Accessed August 2022. Available: <https://www.apha.org/topics-and-issues/climate-change>
- ⁵³ Seervai S., Gustafsson L., Abrams M. 2022. How the US Health Care System Contributes to Climate Change. Commonwealth Fund. Accessed August 2022. Available: <https://www.commonwealthfund.org/publications/explainer/2022/apr/how-us-health-care-system-contributes-climate-change>
- ⁵⁴ Seervai S., Gustafsson L., Abrams M. 2022. How the US Health Care System Contributes to Climate Change. Commonwealth Fund. Accessed August 2022. Available: <https://www.commonwealthfund.org/publications/explainer/2022/apr/how-us-health-care-system-contributes-climate-change>
- ⁵⁵ Salas R., Friend T., Bernstein A., Jha A. 2020. Adding a Climate Lens to Health Policy in the United States. Health Affairs. 39(12). <https://doi.org/10.1377/hlthaff.2020.01352>
- ⁵⁶ Health Care Climate Council. 2020. Climate Action: A Playbook for Hospitals. Accessed August 2022. Available: <https://climatecouncil.noharm.org>
- ⁵⁷ Washington State Department of Commerce. 2022. Weatherization Programs. Accessed August 2022. Available: <https://www.commerce.wa.gov/growing-the-economy/energy/weatherization-and-energy-efficiency/>
- ⁵⁸ ASHRAE. 2020. ASHRAE Position Document on Indoor Air Quality. Accessed August 2022. Available: https://www.ashrae.org/file%20library/about/position%20documents/pd_indoor-air-quality-2020-07-01.pdf

Bree Collaborative Pediatric Asthma Guidelines

Updated: September 13, 2022

⁵⁹ National Center for Healthy Housing. 2022. Asthma Information and Evidence. Accessed August 2022. Available: <https://nchh.org/information-and-evidence/healthy-housing-policy/state-and-local/asthma/>

⁶⁰ Brookings Center for Health Policy. 2015. Community Based Approaches for Optimal Asthma Outcomes and Accountable Population Health. Accessed September 2022. Available: <https://www.aafa.org/media/1535/aafa-brookings-report-2015.pdf>

⁶¹ Farmer S., Patel ., George M., McStay F., Hart R., & McClellan M. 2015. A case study in payment reform to support optimal pediatric asthma care. Brookings Institute. Accessed July 2022. Available: <https://www.brookings.edu/research/a-case-study-in-payment-reform-to-support-optimal-pediatric-asthma-care/>

⁶² Washington State Department of Health. 2020. Asthma Data. Accessed August 2022. Available: <https://doh.wa.gov/data-statistical-reports/diseases-and-chronic-conditions/asthma-data>

⁶³ Centers for Disease Control and Prevention. 2018. Vital Signs: Asthma in Children. Accessed August 2022. Available: <https://www.cdc.gov/vitalsigns/childhood-asthma/index.html>

⁶⁴ WA Department of Health. (2010). Children and Youth with Asthma: Washington State 2009-2010. <https://www.doh.wa.gov/Portals/1/Documents/Pubs/345-305-ChildAsthmaFactSheet.pdf>

⁶⁵ WA Department of Health. (2022). Asthma Data. <https://www.doh.wa.gov/dataandstatisticalreports/diseasesandchronicconditions/asthmadata>