

**Consensus Recommendations for Personal Protective Equipment Use
Collection of Nasopharyngeal (NP) or Oropharyngeal (OP) Swabs and Workplace Exposure**

Developed by the Washington Healthcare Forum Work Group – 3/23/2020

In the face of the COVID-19 outbreak, health care organizations in Washington state are working together to create a common set of high-level recommendations that individual organizations will use to develop their own individual policies and procedures related to the protection of their employees and patients.

These recommendations are predicated on three key principles:

1. Our highest priority is the health and safety of our employees and our patients. We must maintain a stable and healthy workforce as our health care organizations come together to face this public health crisis and continue to serve our patients and communities.
2. These shared recommendations will be based on the best available, most recent scientific evidence. As the science around the COVID-19 virus is rapidly evolving, health care organizations are committed to remaining current and updating these guidelines as needed, based on new findings.
3. These recommendations are designed as a high-level resource for individual organizations. Health care services related to the COVID-19 virus will be delivered in many different settings to many different types of patients. Each enterprise will ultimately have to utilize their best clinical judgment in how they apply and adapt these recommendations in the day-to-day delivery of patient care services.

Collection of Nasopharyngeal (NP) or Oropharyngeal (OP) Swabs

This consensus recommendation relates to collection of nasopharyngeal (NP) or oropharyngeal (OP) swabs. COVID-19 is an emerging disease and our knowledge about the natural history, transmission, clinical course and treatment is constantly updated. When the CDC announced isolation precautions with associated recommendations for personal protective equipment, they differed from the World Health Organization (WHO) by recommending airborne and contact precautions due to this being a new virus. The WHO has continued to recommend droplet and contact precautions, based on SARS-CoV-2 being primarily transmitted through respiratory droplets, reserving airborne precautions for those patients undergoing aerosol-generating procedures. Washington led the change to adopt the WHO guidelines and now Oregon has followed. Similar CDC changes are expected soon.

Aerosol Generating Procedures

A list of aerosol-generating procedures that would still require airborne precautions follows:

- Intubation, extubation and related procedures such as manual ventilation and open
 - suctioning
- Tracheotomy and tracheostomy procedures (insertion, open suctioning, removal)
- Bronchoscopy

- Surgery and post-mortem procedures involving high-speed devices
- Some dental procedures (such as high-speed drilling)
- Non-invasive ventilation (NIV) such as bi-level positive airway pressure (BiPAP) and
 - continuous positive airway pressure ventilation (CPAP)
- High-frequency oscillating ventilation (HFOV)
- High-flow nasal oxygen (HFNO), also called high-flow nasal cannula
- Induction of sputum
- Medication administration via continuous nebulizer

Mechanical ventilation – judged on a case by case basis by the ICU attending with Infectious Disease and/or Infection Prevention

- As long as the ICU ventilator circuit has a filter in place and the circuit remains closed, airborne precautions are not needed
- If patient care requires, or is at risk for, frequent breaks in the circuit, airborne precautions should be used

Collection of nasopharyngeal (NP) or oropharyngeal (OP) swabs

Outpatient collection of NP or OP swabs is not considered an aerosol generating procedure. This was determined in conjunction with Infectious Disease and Infection Prevention providers from around the region.

Providers collecting such samples should use droplet and contact precautions with mask, eye protection, gown and gloves. Collecting nasopharyngeal swabs should be done correctly to maximize the sensitivity and specificity of the test result; and staff education will be provided.

Severely ill patients requiring hospitalization should have testing deferred prior to transfer. Inpatient collection of NP or OP swabs for critically ill or highly symptomatic patients may still require the addition of airborne precautions. The attending physician in conjunction with ID or Infection Prevention can assist with this determination.

Workplace Exposure

Health care workers who experience a low-risk exposure may continue working without a mask if they are asymptomatic. Health care workers may continue to work after a medium- or high-risk exposure if they are (1) asymptomatic and (2) undergo symptom and temperature monitoring at the start of each shift.

The Work Group will continue to update these recommendations as new information becomes available and will be developing future recommendations to further align and coordinate our collective response. Updates include:

3/23/2020 – Added language post high-risk exposure

About the Washington Healthcare Forum

Founded in 1999, the Washington Healthcare Forum is a coalition of physicians, hospitals, health plans, and state associations that have joined to improve the local health care system.

Work group members who developed this policy

Jason Lake, MD, Confluence Health
Tracy Kasnic, R.N., BSN, MBA, CENP, Confluence Health
Jennifer Graves, RN, MS, Kaiser Permanente
Bill Robertson, MultiCare
Dave Carlson, MD, MultiCare
Erica Peavy, MD, PolyClinic
Amy Compton-Phillips, MD, Providence
Mary Cranstoun, CEBS, Providence
Tim Delitt, MD, UW Medicine
Sue Anderson, Virginia Mason
Darcy Jaffe, MSN, ARNP, Washington State Hospital Association
Jessica Martinson, Washington State Medical Association

Board of Directors

Kris Johnson, President and CEO, Association of Washington Business
Chris Bandoli, Executive Director, Association of Washington Healthcare Plans
Peter Rutherford, MD, CEO, Confluence Health
Aric Coffman, MD, CEO and Market President, The Everett Clinic, Part of Optum Care
Jaja Okigwe, President and CEO, First Choice Health Network
Susan Mullaney, President, Kaiser Foundation Health Plan of Washington
Peter Adler, President and Regional Vice President, Molina of Washington
Bill Robertson, President and CEO, MultiCare
Jeff Roe, President and CEO, Premera Blue Cross
Elaine Couture, EVP and CEO, Washington- Montana, Providence St. Joseph Health
Tim Lieb, President, Regence Blue Shield in Washington
Lisa Brandenburg, Chief Health System Officer, UW Medicine
Gary Kaplan, MD, Chairman and CEO, Virginia Mason
Cassie Sauer, President and CEO, Washington State Hospital Association
Jennifer Hanscom, Executive Director and CEO, Washington State Medical Association

Staff

Rick Rubin, Executive Director, Washington Healthcare Forum

OneHealthPort manages the affairs of the Forum. Rick Rubin, OneHealthPort CEO, serves as the Executive Director of the Forum. His contact info is: rickr@onehealthport.com, 206-624-3128 ext. 111.